


Section 12.6: Quadratic Surfaces

Ellipse:

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$



with $a \neq b$



Hyperbola:

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$$

$$-\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$



A second-degree equation in three variables x , y , and z may be expressed in one of two standard forms

$$Ax^2 + By^2 + Cz^2 + E = 0 \quad \text{or} \quad Ax^2 + By^2 + Cz = 0$$

where A, B, C, E are constants. To sketch the graph of a quadratic surface, it is useful to determine the curves of intersection of the surface with planes parallel to the coordinate planes. These curves are called **traces** or **cross-sections** of the surface.

Quadratic surfaces can be grouped into 5 categories: **quadratic cylinders**(cylindrical surfaces from 12.1 notes), **ellipsoids**, **hyperboloids**, **cones**, and **paraboloids**.

For the following examples, assume that $a > 0$, $b > 0$, and $c > 0$.

Ellipsoid:

standard equation: $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$

intercepts: $(\pm a, 0, 0)$, $(0, \pm b, 0)$, and $(0, 0, \pm c)$

cross-sections: (when they exist)


parallel to xy -plane ($z = k$): ellipse

parallel to xz -plane ($y = k$): ellipse

parallel to yz -plane ($x = k$): ellipse

Note: If $a = b = c$ the figure is a sphere. If only two of the constants are equal then the figure is an ellipsoid with the trace involving the two constants being a circle.

Ellipsoid



Pg 4: Hyperboloid(one sheet)

Hyperboloid of one sheet.

standard equation: $\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 1$


cross-sections:

parallel to xy -plane($z = k$): ellipse

parallel to xz -plane($y = k$): hyperbola


parallel to yz -plane($x = k$): hyperbola

Note the axis of the hyperboloid corresponds to the variable whose coefficient is negative.



hyperboloid (one sheet)

Example: Sketch the graph of $x^2 - \frac{y^2}{9} + z^2 = 1$



Let $y=0$: $x^2 + z^2 = 1$

$y = \pm 1$ $x^2 - \frac{1}{9} + z^2 = 1$

$x^2 + z^2 = 1 + \frac{1}{9}$

$y = \pm 2$ $x^2 - \frac{4}{9} + z^2 = 1$

$x^2 + z^2 = 1 + \frac{4}{9}$

Pg 6: hyperboloid (two sheets)

Hyperboloid of two sheets.

standard equation: $-\frac{x^2}{a^2} - \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$


cross-sections:

parallel to xy -plane ($z = k$): ellipse (when they exist)

parallel to xz -plane ($y = k$): hyperbola

parallel to yz -plane ($x = k$): hyperbola

Note the axis of the hyperboloid corresponds to the variable whose coefficient is positive.



Pg 7: cones

Cones:

standard equation: $\frac{x^2}{a^2} + \frac{y^2}{b^2} = \frac{z^2}{c^2}$

or $\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 0$


Note: If $a = b$ then we say we have a circular cone.

cross-sections:

parallel to xy -plane ($z = k$): ellipse

parallel to xz -plane ($y = k$): hyperbola for $k \neq 0$, 2 lines if $k = 0$

parallel to yz -plane ($x = k$): hyperbola for $k \neq 0$, 2 lines if $k = 0$




Example: Sketch the graph of $z^2 = x^2 + y^2$

Circular Cone

$$z=0 \quad x^2 + y^2 = 0$$

$$z=\pm 1 \quad x^2 + y^2 = 1$$



Paraboloids:

Elliptic paraboloid

standard equation:

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = \frac{z}{c}$$

Note: If $a = b$ then we say we have a circular paraboloid.


cross-sections:

parallel to xy -plane ($z = k$): ellipse for $k > 0$

parallel to xz -plane ($y = k$): parabola

parallel to yz -plane ($x = k$): parabola


Note the axis of the paraboloid corresponds to the variable raised to the first power.



$$z = \underline{x^2 + y^2 + 10}$$
$$z = \underline{10 - x^2 - y^2}$$

Example: Sketch the graph of $z = \frac{x^2}{4} + \frac{y^2}{9}$

elliptical paraboloid.



Pg 11: Hyperbolic Paraboloid

hyperbolic paraboloid

standard equation: $\frac{x^2}{a^2} - \frac{y^2}{b^2} = \frac{z}{c}$

cross-sections:

parallel to xy -plane ($z = k$): hyperbola for $k > 0$

parallel to xz -plane ($y = k$): parabola

parallel to yz -plane ($x = k$): parabola

Note the axis of the paraboloid corresponds to the variable raised to the first power.

